
【Review Article】

Step by Step: Accumulated Knowledge and Future Directions of Step-defined Ambulatory Activity

John M. Schuna, Jr.,¹⁾ Catrine Tudor-Locke¹⁾

1) Walking Behavior Laboratory, Pennington Biomedical Research Center

ABSTRACT Walking is one of the most common forms of human locomotion. The development of objective monitoring devices (e.g., pedometers or accelerometers) has afforded public health researchers a unique opportunity to measure ambulatory behavior, including walking, with minimal bias. Objective monitors have been used in a number of cross-sectional national- and state-representative studies to assess walking behavior. Data from these studies, and smaller investigations, have allowed researchers to assess the relationships between total daily ambulation (steps/day) and various health indicators. The growing body of investigations measuring ambulatory behavior has also prompted researchers to seek answers to related questions: “how many steps/day are too few?” and “how many steps/day are enough?” Moreover, recent epidemiological evidence has linked ambulatory speed (steps/min) to various health outcomes and has peaked researchers’ interests in answering “how fast is enough?” This brief review: 1) summarizes current epidemiological literature examining objectively monitored ambulatory behavior, 2) answers public health relevant questions concerning insufficient and sufficient amounts of daily walking, 3) considers the relative importance of walking speed in relation to public health, and 4) identifies future research directions related to the assessment of walking behavior.

Key words: walking, public health, physical activity, sedentary lifestyle, epidemiology

Introduction

Numerous health benefits are associated with a physically active lifestyle.¹⁾ Although there are a myriad of distinct types of physical activity an individual may engage in, walking is the most commonly reported leisure-time physical activity among adults.²⁾ Moreover, walking is an integral part of nearly all forms of daily locomotion and is a natural, simple, accessible, and effective mode of human movement. Considering the ubiquitous nature of walking, its current and future importance to international public health should not be overlooked.

Walking can be most precisely measured using objective monitoring devices (e.g., pedometers or accelerometers) that quantify movement in terms of

steps. Step counts from such devices are often (but not always) provided in real-time (i.e., using a digital read out) and can be used to characterize ambulation at the individual or population level. To aid interpretation of habitual behavior, step counts are typically expressed as steps/day, which provides practitioners and members of the lay public with an easy to understand metric representing the total volume of ambulatory behavior (including walking) in a given day. Guidelines for physical activity from public health authorities have typically prescribed physical activity in terms of frequency, duration, and intensity; however, recent step-based physical activity recommendations have been made by various health groups, professional organizations, and government entities.³⁻⁷⁾

The multiple purposes of this brief article include assembling the rapidly accumulating knowledge concerning the epidemiology of objectively monitored ambulatory activity, summarizing emerging answers to step-defined indices concerning insufficient and

Address for correspondence: Catrine Tudor-Locke;
Walking Behavior Laboratory, Pennington Biomedical
Research Center; 6400 Perkins Road, Baton Rouge, LA,
70808, USA; Tudor-Locke@pbrc.edu

sufficient walking doses, pondering the role of walking speed relative to these questions, and identifying future research needs.

Epidemiology of objectively monitored ambulatory activity

The body of literature focused on step-defined physical activity has grown quickly over the last decade and a clearer understanding of its distribution and determinants is becoming more apparent. A number of national- and state-representative surveys have included objective monitoring of steps/day (by accelerometer or pedometer) and these are assembled in Figure 1,⁸⁻¹⁵⁾ juxtaposed against the highest average values collected to date in an Amish sample.¹⁶⁾ The representative surveys were conducted in Canada,⁹⁾ Japan,¹¹⁾ Norway,¹⁰⁾ Switzerland,¹³⁾ Western Australia,¹²⁾ and in the United States (U.S.) as part of Colorado on the Move,¹⁵⁾ America on the Move,⁸⁾ and the National Health and Nutrition Examination Survey (NHANES).¹⁴⁾ Since Amish people choose to live their lives without modern technologies, including motorized transportation, their pedometer-determined data¹⁶⁾ provide a stark example of what life might have been like before such contemporary advances.

Regardless of the choice of instrumentation (accelerometer or pedometer), it is apparent that daily step counts within the U.S. are the lowest population-level values collected to date.^{8,14,15)} The only country reporting trend data based on multi-year surveillance has been Japan;¹⁷⁾ between 1995 and 2007 there has been a population-level decline in objectively determined physical activity on the order of 529 and 857 steps/day for men and women, respectively, from a peak in 1998-2000. Significantly, the age-adjusted proportion of people taking < 4,000 steps/day has increased by 4.8% in men and 8.2% in women.

With few exceptions,¹⁰⁾ studies of step-defined physical activity indicate that men take more steps/day on average than women.^{8,9,11-14,16)} Steps/day is also inversely related with age¹⁸⁾ and BMI.^{15,19-25)} The number of daily accumulated steps also reflects day of the week, with weekend days being typically lower than weekdays,²⁶⁾ month,²⁷⁾ and season.^{27,28)} For example, Yasunaga et al.²⁹⁾ and Mitsui et al.³⁰⁾ have reported that steps/day are lower in winter months compared with summer months. Steps/day appears to also be impacted by weather changes and perceived neighborhood walkability.³¹⁻³³⁾ In particular, having local and accessible destinations to walk to is associated with higher steps/day.^{34,35)} Engagement in




Figure 1 Mean steps/day among men and women by country, geographic region, or population sub-group.

NHANES = National Health and Nutrition Examination Survey; AOM = America on the Move.

exercise/sports is associated with increased steps/day as is active commuting behaviors.^{26,36,37)} For example, Wener and Evans reported that train commuters averaged 30% more steps/day than car commuters and they were also four times more likely to accumulate 10,000 steps/day.³⁷⁾

How many steps/day are too few?

Nationally representative assessments within the U.S. have revealed that many individuals perform little moderate-to-vigorous physical activity and spend inordinate amounts of time in sedentary pursuits.³⁸⁻⁴⁰⁾ These observations are alarming as high levels of sedentary behavior are associated with increased risks for a variety of negative health outcomes.⁴¹⁻⁴⁴⁾ Such findings provide a basis for asking “how many steps/day are too few?”⁴⁵⁾ That is, below what steps/day threshold are risks for various negative health outcomes significantly increased?

A minimum threshold of 5,000 steps/day was initially suggested by Tudor-Locke and colleagues after observing higher BMI values among adults accumulating < 5,000 steps/day in comparison to those taking > 5,000 steps/day.²²⁾ The authors suggested that taking < 5,000 steps/day may be an indicator of a “sedentary lifestyle index.” Tudor-Locke and Bassett later used 5,000 steps/day as the floor value of their proposed graduated step index.⁴⁶⁾ Subsequently, the 5,000 steps/day threshold has been used by a number of researchers for population surveillance,^{14,17,47-51)} screening,^{52,53)} and intervention.^{52,54)}

In terms of health risks, a number of studies have reported that accumulating < 5,000 steps/day is associated with a higher waist circumference,^{19,24)} body fat percentage,^{20,22-24)} and BMI.^{15,19,20,22-25,55)} Recent investigations have also indicated that the prevalence of hypertension, dyslipidemia, and the metabolic syndrome is higher among individuals taking < 5,000 steps/day in comparison to those accumulating more steps/day.^{56,57)} Additionally, McKercher and colleagues reported that the prevalence of depression was 50% higher among those individuals taking < 5,000 steps/day in comparison to those taking \geq 7,500 steps/day among women, and \geq 12,500 steps/day among men.⁵⁸⁾

Despite the utility of using a straight-forward criterion of < 5,000 steps/day as a marker of ambulatory insufficiency, a certain degree of subjectivity was involved in its development and a range of other daily minimums have been proposed.⁴⁵⁾ Step accumulations of < 6,000 and < 7,500 steps/day have been labeled “inactive” among middle-aged American women and Australian men and women, respectively.^{59,60)} These same thresholds have been labeled “sedentary” in other studies among postmenopausal women.^{61,62)} Another definition, “insufficiently active”, has been defined as < 6,700 steps/day for men and < 5,900 steps/day for women,³¹⁾ while < 4,000 steps/day has been referred to as “sedentary” in several Japanese studies.^{17,63,64)}

Although there is potential usefulness in identifying a singular minimum steps/day for screening and tracking purposes, this approach is imperfect, as relatively lower step counts will likely be associated with increasingly negative outcomes and relatively higher step counts will be associated with better outcomes.⁴⁵⁾ In this sense, there is an inevitable degree of subjectivity in identifying any minimum daily step recommendation in an attempt to answer “how many steps/day are too few?” However, validation studies with longitudinal step data and tracking of various health outcomes may shed light on an appropriate minimum step threshold. Until that time, and considering the substantial body of published literature which has already adopted this threshold, the consistent use of the “sedentary lifestyle index” (< 5,000 steps/day) would allow for comparisons between studies and population groups.⁴⁵⁾

How many steps/day are enough?

Numerous government and agency public health bodies have published guidelines and recommendations which highlight the importance of physical activity and its role in chronic disease prevention.^{3,65-68)} These guidelines have emphasized moderate-to-vigorous physical activity as a primary behavioral target.⁶⁸⁾ In recent times, various step-based recommendations have been put forth.³⁻⁷⁾ Such recommendations should be thought of as a supplement to existing physical activity guidelines, not a

replacement or a source of additional confusion.⁶⁹⁾ Still, it would be useful to identify, in terms of habitual activity levels, “how many steps/day are enough?” while also reflecting existing public health recommendations for frequency, duration, and intensity.

A previous review attempted to answer “how many steps/day are enough?” by computing a steps/day value based upon current physical activity guidelines.⁷⁰⁾ The review referenced previous laboratory studies which identified a cadence of 100 steps/min as a proxy for moderate-to-vigorous physical activity.⁷¹⁻⁷⁵⁾ Using this information, and further assuming that physical activity guidelines typically call for 30 to 60 minutes of moderate-to-vigorous intensity activity per day,^{67,76,77)} a range of 7,100-11,000 steps/day was identified as the daily volume of steps needed to be sufficiently active.⁷⁰⁾ This range of values accounts for a basal minimum of 5,000 steps/day. In order to truly meet physical activity guidelines, a portion of the abovementioned range (3,000 steps/day or 15,000 steps/week) should be completed in bouts of at least 10 minutes and at an intensity of at least 100 steps/min.⁷⁰⁾

Results from a series of studies that reported steps/day translations for time spent in moderate-to-vigorous physical activity and energy expenditure present another option for addressing the “how many steps/days are enough?” question.^{50,78-82)} Findings from these studies were summarized by Tudor-Locke and colleagues who reported that a threshold of 7,000-8,000 steps/day is indicative of an individual meeting minimal physical activity or energy expenditure guidelines.⁷⁰⁾ Interestingly, the minimum steps/day value computed using the previously described assumption-based methods (i.e., 7,100 steps/day) falls within the 7,000 to 8,000 steps/day range. The American College of Sports Medicine (ACSM) included achieving $\geq 7,000$ steps/day in its list of evidence-based recommendations collated as part of its 2011 Position Stand providing guidance for prescribing exercise.⁷⁾

The “how many steps/day are enough?” question can also be evaluated from a health risk perspective. Krumm and colleagues reported that post-menopausal women who took 5,000-7,500 steps/day had sig-

nificantly lower BMI values than those accumulating $< 5,000$ steps/day. Moreover, women who took 7,500-9,999 steps/day had significantly lower BMI values than those accumulating 5,000-7,500 steps/day. Despite this evidence, the general lack of information relating steps/day to health outcomes makes it difficult to answer “how many steps/day are enough?”

How fast is enough?

A measure of steps/day is an indicator of total volume of ambulatory activity, regardless of the speed at which it is accumulated. A meta-analysis of prospective cohort studies indicated that self-reported walking pace (or speed) was a stronger independent predictor of all-cause mortality than self-reported walking volume (48 vs. 26% reductions in risk, respectively).⁸³⁾ Recently, some researchers have begun to capture and interpret cadence (step/min) in an effort to provide more information about free-living walking speed beyond simply using a measure of steps/day.⁸⁴⁻⁸⁸⁾ Walking speed is determined chiefly by cadence and stride length; cadence is the primary strategy employed to increase walking speed, giving over to stride lengthening as an individual transitions from walking to jogging to running.⁸⁹⁾

A number of controlled studies⁷¹⁻⁷⁵⁾ have demonstrated that cadence is strongly related to intensity ($r = 0.93$)⁸⁹⁾ and have consistently concluded that, despite inevitable inter-individual variability, a cadence of 100 steps/min is a reasonable heuristic value indicative of moderate intensity (defined as 3 METs; metabolic equivalents; 1 MET = 3.6 ml of oxygen consumed per kg body weight per minute). Based on eight different covert observation studies⁹⁰⁻⁹⁷⁾ of adults walking in parks, shopping centers, and city streets around the world, the mean weighted observed cadence is 115.2 steps/min,⁸⁹⁾ clearly showing that cadences of > 100 steps/min are easily attainable in natural settings. However, analyses of accelerometer data collected as part of the U.S. NHANES indicates that time accumulated over 100 steps/min is a seemingly rare phenomenon in daily life for the average U.S. citizen.⁸⁸⁾ In fact, these data show that U.S. adults average < 7 min/day at such a cadence but they do spend > 383 min/day at a cadence of 1-19 steps/

min and > 289 min/day at zero cadence (indicative of non-movement).

Faced with the rarity of moderate intensity free-living ambulation, but needing to rank individuals for surveillance and empirical inquiry nonetheless, some researchers have begun to report peak cadence indicators, specifically peak 30-minute cadence, defined as the mean steps/min for the highest (not necessarily consecutive) minutes in a day using the output from time-stamped accelerometers.^{86,87,98)} Peak 30-minute cadence has been described as an indicator of “best natural effort.”⁸⁷⁾ U.S. men average a peak 30-minute cadence of 73.7 steps/min, and despite their generally shorter stature (and therefore relatively higher cadence measured during short distance gait tests⁹⁹⁾, U.S. women have a lower average peak 30-minute cadence of 69.6 steps/min.⁸⁷⁾ Peak 30-minute cadence is also related to age, BMI, and steps/day.^{87,98)}

Conclusions/Future research needs

Walking is integral to daily mobility and is also a part of a healthy lifestyle. Although a number of national- and state-representative assessments of step-defined physical activity have been published, continued collection of data representing other regions will further inform distribution characteristics of this objectively determined indicator of mobility and health. In addition, continued secondary analysis of existing health-related data sets will illuminate important relationships between steps/day and various health indicators. Widespread adoption of systematic on-going surveillance of step-defined physical activity outside of Japan would provide more precise international tracking data.

The available cross-sectional literature suggests that ambulatory volume (steps/day) is inversely associated with waist circumference, body fat percentage, and BMI; however, more longitudinal and intervention studies are needed if dose-response issues (e.g., “how many steps/day are too few?” or “how many steps/day are enough?”) are to be clarified. Such studies may elucidate the relationships between walking and various negative health outcomes (e.g., obesity, heart disease, diabetes, etc.), while evaluating the best approaches to favorably modify walking behavior.

Factors which appear to influence steps/day include climatic forces and perceived neighborhood walkability, however, additional research is required to explore, manipulate, and evaluate other factors that may shape this critical health behavior. Using cadence to capture and study ambulatory intensity in free-living is a novel frontier and more investigations are needed to elucidate the relationships between habitual best cadence measures (e.g., peak 30-minute cadence) and various negative health outcomes. In addition, it has yet to be determined whether or not cadence provides any more useful information compared to only using a simple volume indicator of steps/day. Given the obvious worldwide interest in step-defined physical activity research, it would be beneficial to foment international collaborations in an effort to support and implement this next phase of research.

References

- 1) Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Report, 2008. U.S. Department of Health and Human Services, Washington, DC, 2008.
- 2) Ham SA, Kruger J, Tudor-Locke C. Participation by US adults in sports, exercise, and recreational physical activities. *J Phys Act Health*. 2009; 6(1): 6-14.
- 3) Ishikawa-Takata K, Tabata I. Exercise and physical activity reference for health promotion 2006 (EPAR2006). *J Epidemiol*. 2007; 17(5): 177.
- 4) National Obesity Forum. Useful tools and agencies: Pedometers National Obesity Forum, 2006. <http://www.nationalobesityforum.org.uk/health-care-professionals-mainmenu-155/treatment-mainmenu-169/192-useful-tools-and-agencies.html>: Accessed 08/06/2012
- 5) Queensland Health. 10,000 Steps. Queensland Health, 2012. <http://www.10000steps.org.au>: Accessed 08/08/2012
- 6) The President’s Council on Fitness, Sports & Nutrition. The President’s Challenge: Presidential Active Lifestyle Award (PALA+). The President’s Council on Fitness, Sports & Nutrition, 2012. <https://www.presidentschallenge.org/challenge/>

- active/index.shtml: Accessed 08/08/2012
- 7) Garber CE, Blissmer B, Deschenes MR, et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011; 43(7): 1334-59.
 - 8) Bassett DR, Jr, Wyatt HR, Thompson H, Peters JC, Hill JO. Pedometer-measured physical activity and health behaviors in United States adults. *Med Sci Sports Exerc.* 2010; 42(10): 1819-25.
 - 9) Colley RC, Garriguet D, Janssen K, Craig CL, Clarke J, Tremblay MS. Physical activity of Canadian adults: Accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health Rep.* 2011; 22(1): 1-8.
 - 10) Hansen BH, Kolle E, Dyrstad SM, Holme I, Anderssen SA. Accelerometer-determined physical activity in adults and older people. *Med Sci Sports Exerc.* 2012; 44(2): 266-72.
 - 11) Inoue S, Ohya Y, Odagiri Y, et al. Sociodemographic determinants of pedometer-determined physical activity among Japanese adults. *Am J Prev Med.* 2011; 40(5): 566-71.
 - 12) McCormack G, Giles-Corti B, Milligan R. Demographic and individual correlates of achieving 10,000 steps/day: Use of pedometers in a population-based study. *Health Promot J Austr.* 2006; 17(1): 43-7.
 - 13) Sequeira MM, Rickenbach M, Wietlisbach V, Tullen B, Schutz Y. Physical activity assessment using a pedometer and its comparison with a questionnaire in a large population survey. *Am J Epidemiol.* 1995; 142(9): 989-99.
 - 14) Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps per day in US adults. *Med Sci Sports Exerc.* 2009; 41(7): 1384-91.
 - 15) Wyatt HR, Peters JC, Reed GW, Barry M, Hill JO. A Colorado statewide survey of walking and its relation to excessive weight. *Med Sci Sports Exerc.* 2005; 37(5): 724-30.
 - 16) Bassett DR, Jr, Schneider PL, Huntington GE. Physical activity in an Old Order Amish community. *Med Sci Sports Exerc.* 2004; 36(1): 79-85.
 - 17) Inoue S, Ohya Y, Tudor-Locke C, Tanaka S, Yoshiike N, Shimomitsu T. Time trends for step-determined physical activity among Japanese adults. *Med Sci Sports Exerc.* 2011; 43(10): 1913-9.
 - 18) Tudor-Locke C, Hart TL, Washington TL. Expected values for pedometer-determined physical activity in older populations. *Int J Behav Nutr Phys Act.* 2009; 6: 59.
 - 19) Dwyer T, Hosmer D, Hosmer T, et al. The inverse relationship between number of steps per day and obesity in a population-based sample: the AusDiab study. *Int J Obes (Lond).* 2007; 31(5): 797-804.
 - 20) Krumm EM, Dessieux OL, Andrews P, Thompson DL. The relationship between daily steps and body composition in postmenopausal women. *J Womens Health.* 2006; 15(2): 202-10.
 - 21) Tudor-Locke C, Bassett DR, Jr, Rutherford WJ, et al. BMI-referenced cut points for pedometer-determined steps per day in adults. *J Phys Act Health.* 2008; 5(Suppl 1): 126-39.
 - 22) Tudor-Locke C, Ainsworth BE, Whitt MC, Thompson RW, Addy CL, Jones DA. The relationship between pedometer-determined ambulatory activity and body composition variables. *Int J Obes Relat Metab Disord.* 2001; 25(11): 1571-8.
 - 23) Mitsui T, Shimaoka K, Tsuzuku S, Kajiooka T, Sakakibara H. Pedometer-determined physical activity and indicators of health in Japanese adults. *J Physiol Anthropol.* 2008; 27(4): 179-84.
 - 24) Hornbuckle LM, Bassett DR, Jr, Thompson DL. Pedometer-determined walking and body composition variables in African-American women. *Med Sci Sports Exerc.* 2005; 37(6): 1069-74.
 - 25) Cook I, Alberts M, Lambert EV. Relationship between adiposity and pedometer-assessed ambulatory activity in adult, rural African women. *Int J Obes (Lond).* 2008; 32(8): 1327-30.
 - 26) Tudor-Locke C, Jones R, Myers AM, Paterson DH, Ecclestone NA. Contribution of structured exercise class participation and informal walking for exercise to daily physical activity in community-dwelling older adults. *Res Q Exerc Sport.* 2002; 73(3): 350-6.

- 27) Kang M, Basset DR, Barreira TV, Tudor-Locke C, Ainsworth BE. Measurement effects of seasonal and monthly variability on pedometer-determined data. *J Phys Act Health*. 2012; 9(3): 336-43.
- 28) Dasgupta K, Chan C, Da Costa D, et al. Walking behaviour and glycemic control in type 2 diabetes: Seasonal and gender differences - study design and methods. *Cardiovasc Diabetol*. 2007; 6: 1.
- 29) Yasunaga A, Togo F, Watanabe E, Park H, Shephard RJ, Aoyagi Y. Sex, age, season, and habitual physical activity of older Japanese: The Nakanojo study. *J Aging Phys Act*. 2008; 16(1): 3-13.
- 30) Mitsui T, Barajima T, Kanachi M, Shimaoka K. Daily walking activity among male office workers in a rural town in northern Japan. *J Physiol Anthropol*. 2010; 29(1): 43-6.
- 31) Oka K, Shibata A. Determinants of meeting the public health recommendations for physical activity among community-dwelling elderly Japanese. *Curr Aging Sci*. 2012; 5(1): 58-65.
- 32) Chan CB, Ryan DA, Tudor-Locke C. Relationship between objective measures of physical activity and weather: A longitudinal study. *Int J Behav Nutr Phys Act*. 2006; 3: 21.
- 33) Van Dyck D, Deforche B, Cardon G, De Bourdeaudhuij I. Neighborhood walkability and its particular importance for adults with a preference for passive transport. *Health Place*. 2009; 15(2): 496-504.
- 34) Kondo K, Lee JS, Kawakubo K, et al. Association between daily physical activity and neighborhood environments. *Environ Health Prev Med*. 2009; 14(3): 196-206.
- 35) King WC, Belle SH, Brach JS, Simkin-Silverman LR, Soska T, Kriska AM. Objective measures of neighborhood environment and physical activity in older women. *Am J Prev Med*. 2005; 28(5): 461-9.
- 36) Tudor-Locke C, Basset DR, Swartz AM, et al. A preliminary study of one year of pedometer self-monitoring. *Ann Behav Med*. 2004; 28(3): 158-62.
- 37) Wener RE, Evans GW. A morning stroll: Levels of physical activity in car and mass transit commuting. *Environ Behav*. 2007; 39(1): 62-74.
- 38) Matthews CE, Chen KY, Freedson PS, et al. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol*. 2008; 167(7): 875-81.
- 39) Troiano RP, Berrigan D, Dodd KW, Mâsse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008; 40(1): 181-8.
- 40) Tucker JM, Welk GJ, Beyler NK. Physical activity in U.S. adults: Compliance with the Physical Activity Guidelines for Americans. *Am J Prev Med*. 2011; 40(4): 454-61.
- 41) Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc*. 2009; 41(5): 998-1005.
- 42) Hu FB, Li TY, Colditz GA, Willett WC, Manson JE. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *JAMA*. 2003; 289(14): 1785-91.
- 43) Ford ES, Kohl HW, III, Mokdad AH, Ajani UA. Sedentary behavior, physical activity, and the metabolic syndrome among U.S. adults. *Obes Res*. 2005; 13(3): 608-14.
- 44) Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW. Coronary heart-disease and physical activity of work. *Lancet*. 1953; 265(6796): 1111-20.
- 45) Tudor-Locke C, Craig CL, Thyfault JP, Spence JC. A step-defined sedentary lifestyle index: < 5,000 steps/day. *Appl Physiol Nutr Metab*. In press.
- 46) Tudor-Locke C, Bassett DR, Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med*. 2004; 34(1): 1-8.
- 47) Sisson SB, Camhi SM, Church TS, Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps/day and metabolic syndrome. *Am J Prev Med*. 2010; 38(6): 575-82.
- 48) Sisson SB, Camhi SM, Tudor-Locke C, Johnson WD, Katzmarzyk PT. Characteristics of step-defined physical activity categories in U.S. adults.

- Am J Health Promot. 2012; 26(3): 152-9.
- 49) Tudor-Locke C, Brashear MM, Johnson WD, Katzmarzyk PT. Accelerometer profiles of physical activity and inactivity in normal weight, overweight, and obese U.S. men and women. *Int J Behav Nutr Phys Act.* 2010; 7: 60.
 - 50) Tudor-Locke C, Johnson WD, Katzmarzyk PT. Relationship between accelerometer-determined steps/day and other accelerometer outputs in U.S. adults. *J Phys Act Health.* 2011; 8(3): 410-9.
 - 51) Tudor-Locke C, Leonardi C, Johnson WD, Katzmarzyk PT, Church TS. Accelerometer steps /day translation of moderate-to-vigorous activity. *Prev Med.* 2011; 53(1-2): 31-3.
 - 52) Bell GJ, Harber V, Murray T, Courneya KS, Rodgers W. A comparison of fitness training to a pedometer-based walking program matched for total energy cost. *J Phys Act Health.* 2010; 7(2): 203-13.
 - 53) Musto A, Jacoba K, Nash M, DelRossi G, Perry A. The effects of an incremental approach to 10,000 steps/day on metabolic syndrome components in sedentary overweight women. *J Phys Act Health.* 2010; 7(6): 737-45.
 - 54) Villanova N, Pasgui F, Burzacchini S, et al. A physical activity program to reinforce weight maintenance following a behavior program in overweight/obese subjects. *Int J Obes (Lond).* 2006; 30(4): 697-703.
 - 55) Inoue S, Ohya Y, Tudor-Locke C, Yoshiike N, Shimomitsu T. Step-defined physical activity and cardiovascular risk among middle-aged Japanese: The National Health and Nutrition Survey of Japan 2006. *J Phys Act Health.* In press.
 - 56) Schmidt MD, Cleland VJ, Shaw K, Dwyer T, Venn AJ. Cardiometabolic risk in younger and older adults across an index of ambulatory activity. *Am J Prev Med.* 2009; 37(4): 278-84.
 - 57) Sisson SB, Camhi SM, Church TS, Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps/day and metabolic syndrome. *Am J Prev Med.* 2010; 38(6): 575-82.
 - 58) McKercher CM, Schmidt MD, Sanderson KA, Patton GC, Dwyer T, Venn AJ. Physical activity and depression in young adults. *Am J Prev Med.* 2009; 36(2): 161-4.
 - 59) Thompson DL, Rakow J, Perdue SM. Relationship between accumulated walking and body composition in middle-aged women. *Med Sci Sports Exerc.* 2004; 36(5): 911-4.
 - 60) Tudor-Locke C, Giles-Corti B, Knuiaman M, McCormack G. Tracking of pedometer-determined physical activity in adults who relocate: results from RESIDE. *Int J Behav Nutr Phys Act.* 2008; 5: 39.
 - 61) Lara S, Casanova G, Spritzer PM. Influence of habitual physical activity on body composition, fat distribution and metabolic variables in early postmenopausal women receiving hormonal therapy. *Eur J Obstet Gynecol Reprod Biol.* 2010; 150(1): 52-6.
 - 62) Barbat-Artigas S, Plouffe S, Dupontgand S, Aubertin-Leheudre M. Is functional capacity related to the daily amount of steps in postmenopausal women? *Menopause.* 2012; 19(5): 541-8.
 - 63) Ishikawa-Takata K, Tanaka H, Nanbu K, Ohta T. Beneficial effect of physical activity on blood pressure and blood glucose among Japanese male workers. *Diabetes Res Clin Pract.* 2010; 87(3): 394-400.
 - 64) Park H, Togo F, Watanabe E, et al. Relationship of bone health to yearlong physical activity in older Japanese adults: Cross-sectional data from the Nakanajo Study. *Osteoporosis Int.* 2007; 18(3): 285-93.
 - 65) Commonwealth Department of Health and Aged Care. National physical activity guidelines for Australians. Commonwealth Department of Health and Aged Care, Canberra, AUS, 1999.
 - 66) Department of Health. Strategy statement on physical activity. Department of Health, London, UK, 1996.
 - 67) US Department of Health and Human Services. Physical activity and health: A report of the Surgeon General. Centers for Disease Control and Prevention, Atlanta, US, 1996.
 - 68) US Department of Health and Human Services. 2008 Physical activity guidelines for Americans. US Department of Health and Human Services, Washington, DC, 2008.
 - 69) Tudor-Locke C. Steps to better cardiovascular health: How many steps does it take to achieve

- good health and how confident are we in this number? *Curr Cardiovasc Risk Rep.* 2010; 4(4): 271-6.
- 70) Tudor-Locke C, Craig CL, Brown WJ, et al. How many steps/day are enough? For adults. *Int J Behav Nutr Phys Act.* 2011; 8: 79.
- 71) Abel M, Hannon J, Mullineaux D, Beighle A. Determination of step rate thresholds corresponding to physical activity classifications in adults. *J Phys Act Health.* 2011; 8(1): 45-51.
- 72) Beets MW, Agiovlasis S, Fahs CA, Ranadive SM, Fernhall B. Adjusting step count recommendations for anthropometric variations in leg length. *J Sci Med Sport.* 2010; 13(5): 509-12.
- 73) Marshall SJ, Levy SS, Tudor-Locke CE, et al. Translating physical activity recommendations into a pedometer-based step goal: 3000 steps in 30 minutes. *Am J Prev Med.* 2009; 36(5): 410-5.
- 74) Rowe DA, Welk GJ, Heil DP, et al. Stride rate recommendations for moderate intensity walking. *Med Sci Sports Exerc.* 2011; 43(2): 312-8.
- 75) Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Can J Appl Physiol.* 2005; 30(6): 666-76.
- 76) Public Health Agency of Canada and the Canadian Society for Exercise Physiology. Canada's physical activity guide to healthy active living. Public Health Agency of Canada, Ottawa, ON, 1998.
- 77) Institute of Medicine. Dietary reference intakes for energy, carbohydrates, fiber, fat, protein and amino acids (macronutrients). National Academy of Sciences, Washington, DC, 2002.
- 78) Tudor-Locke C, Ainsworth BE, Thompson RW, Matthews CE. Comparison of pedometer and accelerometer measures of free-living physical activity. *Med Sci Sports Exerc.* 2002; 32(12): 2045-51.
- 79) Miller R, Brown W. Meeting physical activity guidelines and average daily steps in a working population. *J Phys Act Health.* 2004; 1(3): 218-26.
- 80) Behrens TK, Hawkins SB, Dinger MK. Relationship between objectively measured steps and time spent in physical activity among free-living college students. *Meas Phys Educ Exerc Sci.* 2005; 9(2): 67-77.
- 81) Jordan AN, Jurca GM, Locke CT, Church TS, Blair SN. Pedometer indices for weekly physical activity recommendations in postmenopausal women. *Med Sci Sports Exerc.* 2005; 37(9): 1627-32.
- 82) Macfarlane DJ, Chan D, Chan KL, Ho EY, Lee CC. Using three objective criteria to examine pedometer guidelines for free-living individuals. *Eur J Appl Physiol.* 2008; 104(3): 435-44.
- 83) Hamer M, Chida Y. Walking and primary prevention: A meta-analysis of prospective cohort studies. *Br J Sports Med.* 2008; 42(4): 238-43.
- 84) Ayabe M, Aoki J, Kumahara H, Tanaka H. Assessment of minute-by-minute stepping rate of physical activity under free-living conditions in female adults. *Gait Posture.* 2011; 34(2): 292-4.
- 85) Ayabe M, Aoki J, Kumahara H, et al. Minute-by-minute stepping rate of daily physical activity in normal and overweight/obese adults. *Obes Res Clin Pract.* 2011; 5(2): e151-6.
- 86) Gardner AW, Montgomery PS, Scott KJ, Afaq A, Blevins SM. Patterns of ambulatory activity in subjects with and without intermittent claudication. *J Vasc Surg.* 2007; 46(6): 1208-14.
- 87) Tudor-Locke C, Brashear MM, Katzmarzyk PT, Johnson WD. Peak stepping cadence in free-living adults: 2005-2006 NHANES. *J Phys Act Health.* In press.
- 88) Tudor-Locke C, Camhi SM, Leonardi C, et al. Patterns of adult stepping cadence in the 2005-2006 NHANES. *Prev Med.* 2011; 53(3): 178-81.
- 89) Tudor-Locke C, Rowe DA. Using cadence to study free-living ambulatory behavior. *Sports Med.* 2012; 42(5): 381-98.
- 90) Drillis R. Objective recording and biomechanics of pathological gait. *Ann N Y Acad Sci.* 1958; 74(1): 86-109.
- 91) Molen NH, Rozendal RH. Some factors of human gait. *Proc K Ned Akad Wet C.* 1966; 69(4): 522-7.
- 92) Finley FR, Cody KA. Locomotive characteristics of urban pedestrians. *Arch Phys Med Rehabil.* 1970; 51(7): 423-6.

- 93) Molen NH, Rozendal RH, Boon W. Fundamental characteristics of human gait in relation to sex and location. *Proc K Ned Akad Wet C.* 1972; 45(3): 215-23.
- 94) Eke-Okoro ST, Sandlund B. The effects of load, shoes, sex, and direction on the gait characteristics of street pedestrians. *J Hum Movement Stud.* 1984; 10: 107-14.
- 95) Sato H, Ishizu K. Gait patterns of Japanese pedestrians. *J Hum Ergol (Tokyo).* 1990; 19(1): 13-22.
- 96) Sato H, Sako H, Mukae H, Sato A, Takahashi T. Gait patterns of young Japanese women. *J Hum Ergol (Tokyo).* 1991; 20(1): 85-8.
- 97) Hangland A, Cimbalo RS. Human ethology: Age and sex differences in mall walking. *Percept Mot Skills.* 1997; 85(3 Pt 1): 845-6.
- 98) Barreira TV, Katzmarzyk PT, Johnson WD, Tudor-Locke C. Cadence patterns and peak cadence in U.S. children and adolescents: NHANES, 2005-2006. *Med Sci Sports Exerc.* In press.
- 99) Hollman JH, McDade EM, Petersen RC. Normative spatiotemporal gait parameters in older adults. *Gait Posture.* 2011; 34(1): 111-8.

【要旨日本語訳】

ステップ・バイ・ステップ： 歩数に関するこれまでの知見と今後の方向性

歩行は人間にとって最も一般的な運動の形態である。身体活動を客観的に測定する機器（歩数計や加速度計）の開発によって、公衆衛生研究者は歩行などの行動を、最小限のバイアスで測定することができるようになった。これらの機器を活用した国の調査、州の調査なども多い。これらの調査の結果や、小規模な調査の結果によって、1日の総移動量（歩/日）とさまざまな健康指標との関連が検討されている。これらの多くの研究はまた、「何歩くらいだと活動量が少なすぎるのか」あるいは「何歩くらい歩けば十分なのか」といった疑問を引き起こした。更にいうと、最近の疫学研究は、歩くスピード（歩/分）とさまざまな健康指標との関連を示しており、「どのくらいの速さで歩けばよいのか」といった疑問を提示した。本稿では、1) 客観的な測定装置（歩数計、加速度計）を用いた記述疫学研究の結果を要約し、2) どのくらいの歩行量が不十分、あるいは十分なのかに答え、3) 公衆衛生的視点からみた歩行スピードの重要性を考察し、4) 歩行評価に関する今後の研究の方向性を明らかにする。

（この日本語訳は、読者の利便性を考慮して著者の許可のもとに編集委員会が作成したもので、論文の一部ではありません。日本語訳が著者の意図にあっていない可能性がありますので、正確な意味を確認するためには原文をご確認ください）